



# Brave men and timid women? A review of the gender differences in fear and anxiety

Carmen P. McLean<sup>a,\*</sup>, Emily R. Anderson<sup>b</sup>

<sup>a</sup> National Center for PTSD, Boston VA Healthcare System, 150 S. Huntington Ave. (116B-5), Boston, MA 02130, USA

<sup>b</sup> Yale University Child Study Center, USA

## ARTICLE INFO

### Article history:

Received 6 June 2008

Received in revised form 27 May 2009

Accepted 30 May 2009

### Keywords:

Gender

Anxiety

Gender socialization

## ABSTRACT

Substantial evidence indicates that women report greater fear and are more likely to develop anxiety disorders than men. Women's greater vulnerability for anxiety disorders can be partly understood by examining gender differences in the etiological factors known to contribute to anxiety. This review examines evidence for gender differences across a broad range of relevant factors, including biological influences, temperamental factors, stress and trauma, cognitive factors, and environmental factors. Gender differences are observed with increasing consistency as the scope of analysis broadens to molar levels of functioning. Socialization processes cultivate and promote processes related to anxiety, and moderate gender differences across levels of analysis.

© 2009 Elsevier Ltd. All rights reserved.

## Contents

1.	Evidence of gender differences. . . . .	497
1.1.	Gender differences in fear . . . . .	497
1.2.	Gender differences in anxiety disorders . . . . .	497
1.3.	Gender differences within anxiety disorders . . . . .	498
2.	Overview of etiological factors. . . . .	498
2.1.	Biological factors. . . . .	498
2.1.1.	Genetic factors . . . . .	498
2.1.2.	Physiological reactivity. . . . .	498
2.1.3.	Hormonal influences. . . . .	499
2.1.4.	Evolutionary influences . . . . .	499
2.2.	Broad vulnerability factors: higher order factors . . . . .	499
2.2.1.	Negative affectivity . . . . .	499
2.2.2.	Trait anxiety . . . . .	499
2.3.	Broad vulnerability factors: disorder specific factors . . . . .	500
2.3.1.	Anxiety sensitivity. . . . .	500
2.3.2.	Disgust sensitivity. . . . .	500
2.4.	Stress and trauma . . . . .	500
2.4.1.	Gender-specific trauma . . . . .	500
2.4.2.	Resiliency. . . . .	500
2.4.3.	Appraisals of threat . . . . .	500
2.4.4.	Uncontrollability . . . . .	501
2.4.5.	Self-efficacy. . . . .	501
2.4.6.	Worry . . . . .	501
2.4.7.	Rumination . . . . .	501
2.4.8.	Heightened sensitivity to social cues . . . . .	501
2.5.	Behavioral avoidance. . . . .	501

\* Corresponding author.

E-mail address: [carmen.mclean@va.gov](mailto:carmen.mclean@va.gov) (C.P. McLean).

2.6.	Environmental influences . . . . .	502
2.6.1.	Sociocultural influences . . . . .	502
2.6.2.	Gender role socialization . . . . .	502
3.	Conclusion. . . . .	502
	References . . . . .	503

Based on established findings (e.g., Bourdon et al., 1988; Weissman & Merikangas, 1986), most well-articulated research hypotheses predict that women report and/or demonstrate greater fear and anxiety than men. However, the rationale for this expected finding is often absent or refers only to known gender disparities in prevalence rates of anxiety disorders. This circular reasoning highlights the need to identify complex processes that underlie gender differences in anxiety. By delineating the mechanisms that promote and maintain fear and anxiety among men and women, prevention and treatment efforts may be improved.

The purpose of the present paper is to review evidence of gender<sup>1</sup> differences in anxiety and fear. Results of a comprehensive literature search using electronic databases (PsycInfo, Pubmed) are reviewed from a biosystemic model in which gender-specific vulnerabilities at molecular levels of analysis gradually evolve into fully articulated traits through complex, bidirectional interactions with molar level factors. Within this model, men and women are identified as complex systems comprised of various interrelating mechanisms. For heuristic purposes, these mechanisms are organized into several levels of functioning arranged along a continuum of complexity. Processes occurring at any given level are presumed to influence functioning at all other levels. Thus, it is hypothesized that individual differences in etiological factors for anxiety and fear are moderated by socialization processes that prescribe gender-specific expectations for expression of anxiety and the acceptable means of coping with anxiety. These socialization factors influence expression of traits by shaping patterns of reinforcement that cultivate and promote processes related to anxiety.

First, we provide a brief overview of prevalence of reported fear and anxiety disorders across gender, including developmental patterns that likely contribute to these differences. Next, we review gender differences in etiological factors beginning with biological factors including genetic factors, physiological reactivity, hormonal factors, and evolutionary influences. Broad temperamental factors are examined next, including higher order factors, such as negative affectivity, and disorder specific factors, such as anxiety sensitivity and disgust sensitivity. Moving toward more molar levels of analysis, we review gender differences in stress and trauma-related factors, and then examine the cognitive processes associated with anxiety, including appraisals of threat, uncontrollability, worry and rumination, and sensitivity to social cues. Last, we review gender differences in relevant environmental factors, including sociocultural and modeling factors.

## 1. Evidence of gender differences

### 1.1. Gender differences in fear

Gender differences in fear reporting are evident in children as young as 9–12 years of age (Ollendick, 1983). Among adults, a number of self-report studies indicate that women endorse a greater number and severity of fears than men (e.g., Arrindell, 2000; Bourdon et al., 1988; Davey, 1994). For example, Tucker and Bond (1997) found significant gender differences in fear of repulsive animals (e.g.,

snakes) and fear of harmless animals (e.g., dogs), but no differences in fears of bodily injury, social fears, noise fears, or other classic phobias (e.g. enclosed spaces). Consistent with this result, large scale cross-cultural studies have found that *gender disparities are greatest* in fears of harmless animals (Arrindell, Eisemann, Richter, Oei, & Caballo, 2003) and disgust-relevant animals (Davey, McDonald, & Hirisave, 1998) than for fear-relevant animals (e.g., lions, sharks). In contrast, few gender differences have been found for public speaking or social-evaluative fears (Klorman, 1974), possibly due to socialization influences discussed later in this review.

### 1.2. Gender differences in anxiety disorders

Among children, girls are more likely than boys to experience any anxiety disorder (Anderson, Williams, McGee, & Silva, 1987). For example, in a community sample of 1079 adolescents, Lewinsohn, Gotlib, Lewinsohn, Seeley, and Allen (1998), found that girls were more likely than boys to have a current or lifetime diagnosis of an anxiety disorder. Early childhood appears to mark a period of increasing vulnerability to excessive anxiety among girls. By age six, girls are twice as likely as boys to have experienced an anxiety disorder (Anderson et al., 1987; Lewinsohn et al., 1998), a divergence that appears to continue throughout adolescence (Muris & Ollendick, 2002). Adolescent girls report a greater number of worries, more separation anxiety (Campbell & Rapee, 1994; Costello, Egger, & Angold, 2003; Poulton, Milne, Craske, & Menzies, 2001), and are six times more likely to develop generalized anxiety disorder (GAD) than adolescent boys (Bowen, Offord, & Boyle, 1990; McGee et al., 1990). Childhood obsessive compulsive disorder (OCD) is nearly three times as common among boys (Castle, Deale, & Marks, 1995; Karno, Golding, Sorenson, & Burman, 1988) until puberty, at which time the gender differences equalize and then reverse (Pigott, 1998; Rasmussen & Eisen, 1990).

Similar evidence for gender differences exists in adult samples. Large community-based, epidemiological studies show that women are at greater risk than men for most anxiety disorders (Angst & Dobler-Mikola, 1985; Bruce, Yonkers, Otto, Eisen, & Weisberg, 2005). According to lifetime prevalence data from the National Comorbidity Survey, women are more likely than men to have panic disorder, agoraphobia, GAD (Kessler, McGonagle, Zhao, Nelson, & Hughes, 1994), and posttraumatic stress disorder (PTSD) (Kessler, Sonnega, Bromet, Hughes, & Nelson, 1995). Women are approximately twice as likely to meet criteria for certain phobias (Lindal & Stefánsson, 1993), with the pattern of gender differences mirroring research on fear reporting (Himle, McPhee, Cameron, & Curtis, 1989). Gender differences appear to be the least pronounced for social anxiety disorder (SAD) and OCD (Breslau, Chilcoat, Peterson, & Schultz, 2000; Karno & Golding, 1991).

Findings among clinical samples generally parallel community-based samples. For example, a large study conducted by the World Health Organization in 15 countries found that women were consistently more likely than men to present for treatment of panic disorder with agoraphobia (Gater et al., 1998). In contrast to this general parallel between clinical and community samples, men and women with SAD appear equally likely to seek treatment despite evidence from research among community samples suggesting that women are slightly more likely to have social anxiety disorder (Katzelnick & Greist, 2001).

<sup>1</sup> Throughout this review, the term *gender* will be used to refer to both sex and gender, except in cases where measures of gender role are included to control for the sociocultural and psychological processes inherent to the construct of gender.

### 1.3. Gender differences within anxiety disorders

In addition to gender differences across anxiety disorders, there are also differences within anxiety disorders in terms of chronicity, symptomology, and patterns of comorbidity. Panic attacks are experienced at equal rates among men and women (Telch, Lucas, & Nelson, 1989), but certain symptoms of panic attacks, such as shortness of breath, faintness, and smothering sensations are more commonly described as primary panic symptoms among women than men (Sheikh, Leskin, & Klein, 2002). In terms of course, panic disorder is more chronic among women than men (Hollifield et al., 1997; Yonkers et al., 1998) and is associated with higher rates of comorbidity with GAD, somatization disorder, and agoraphobia (Pigott, 1999; Turgeon, Marchand, & Dupuis, 1998). However, the most striking gender difference is observed across levels of agoraphobic avoidance, in which women are disproportionately represented within the highest levels of agoraphobia (Cameron & Hill, 1989; Turgeon et al., 1998).

Although there are some inconsistencies in the literature, women with OCD appear more likely to exhibit cleaning compulsions and aggressive obsessions, while men more commonly exhibit obsessive slowness, symmetry obsessions/compulsions, touching rituals, and sexual or “odd” symptoms (Castle et al., 1995). Further, men who are affected by OCD tend to have an earlier onset and exhibit a chronic, rather than episodic course (Castle et al., 1995; Pigott, 1998).

Research has found that the rates for pure anxiety disorders (i.e. no comorbid diagnoses) are similar across men and women (Ochoa, Beck, & Steer, 1992). Yet, among individuals with anxiety disorders, comorbid substance use disorders are more common among men (Cox, Swinson, & Shulman, 1993), whereas comorbid mood disorders are more common among women (Ochoa et al., 1992; Scheibe & Albus, 1992). Another study demonstrated higher rates of comorbidity and severity among females in the hoarding sub-type of OCD (Wheaton, Timpano, LaSalle-Ricci, & Murphy, 2008).

## 2. Overview of etiological factors

Women's greater vulnerability for anxiety can be partly understood by examining etiological factors related to anxiety across levels of analysis. Beginning with an examination of biological factors, the scope of analysis in this review increases at each level. Although divisions between levels are somewhat arbitrary, they are heuristically useful for purposes of investigation and review.

### 2.1. Biological factors

#### 2.1.1. Genetic factors

A large body of evidence supports a substantial nonspecific genetic contribution to anxiety disorders (Barlow, 2000). Within specific phobias, genetic factors are estimated to account for up to one third of the total variance (Kendler, Neale, Kessler, Heath, & Eaves, 1992) with the greatest contribution among animal phobias (see Muris & Merckelbach, 2001). Panic disorder also has been found to run in families (Goldstein, Wickramaratne, Horwath, & Weissman, 1997) and has demonstrated a high degree of heritability in twin studies (Perna, Caldirola, Arancio, & Bellodi, 1997). The heritability of panic disorder has been hypothesized to reflect the genetic transmission of physiologic risk factors, such as carbon dioxide hypersensitivity (Perna, Bertani, Caldirola, & Bellodi, 1996).

Genetic factors have been shown to relate with anxiety sensitivity (Stein, Jang, & Livesly, 2002) and with other broad vulnerability factors such as neuroticism. In fact, twin studies show that approximately 50% of the variance in neuroticism is accounted for by genetic factors (Lake, Eaves, Maes, Heath, & Martin, 2000). Heritability of these vulnerability factors helps to explain findings that indicate a genetic influence among anxiety disorders which are

clearly connected to environmental factors. For example, higher concordance ratios are found for PTSD among monozygotic than dizygotic twins (Skre, Onstad, Torgersen, Lygren, & Kringlen, 1993), even when controlling for type of trauma exposure (True et al., 1993). Relevant to our understanding of gender differences are data showing that the heritability of anxiety-related vulnerability factors is greater among women than men. Lake and colleagues (2000) found that genetic factors had a significantly greater influence on the individual variability in neuroticism among women than men. Similarly, research with child and adolescent samples consistently shows higher heritability estimates for fear and phobias for girls than boys (Eley, 2001). Such evidence suggests that gender differences in anxiety may largely correspond with a differential genetic predisposition to anxiety disorders or their related vulnerability factors.

Alternatively, weaker heritability estimates of anxiety-related vulnerability factors among men could signify that processes at higher levels of analysis counteract inherited tendencies among boys. Thus, genetic vulnerabilities may be more supported by the social/environmental influences experienced by girls. This hypothesis is consistent with the results of a large twin study recently conducted by Hettema, Prescott, Myers, Neale, and Kendler (2005). In a sample of over 5000 participants, structural equation modeling was used to examine the underlying causes of comorbidity within anxiety disorders. Results showed that despite a nearly two-fold higher prevalence among women, genetic and environmental risk factors for anxiety disorders were similar across gender. Thus, while the types of genetic factors that place individuals at risk for experiencing anxiety disorders may be the same across gender, the relative impact of genetic factors may be greater among women.

#### 2.1.2. Physiological reactivity

The hypothesis that women are more physiologically predisposed to experience anxiety has not been consistently supported. Studies examining skin conductance within conditioning and extinction paradigms have found greater reactivity among men than women: men display higher levels of responding during classical conditioning tasks (Graham, Cohen, & Schmavonian, 1966), slower rates of habituation to a nocuous tone (Maltzman, 1979), and greater responding across vicarious conditioning tasks (Craig & Lowery, 1969).

In contrast to findings from conditioning paradigms, research examining physiological reactivity in response to acute behavioral stress has yielded different results. Several studies have found that men and women show similar levels of reactivity, as measured by skin conductance in response to fearful stimuli (Katkin & Hoffman, 1976), heart rate during social stress challenge (Kelly, Tyrka, Anderson, Price, & Carpenter, 2008), electrodermal reactivity, heart rate during CO<sub>2</sub> inhalation (Kelly, Forsyth, & Karekla (2006) M.M. Kelly, J.P. Forsyth and M. Karekla, Sex differences in response to a panicogenic biological challenge procedure: An experimental evaluation of panic vulnerability in a non-clinical sample, *Behaviour Research and Therapy* 44 (2006), pp. 1421–1430. Article | PDF (198 K) | View Record in Scopus | Cited By in Scopus (6) Kelly et al., 2006), and heart rate during oral examination (Rohrmann, Netter, Hennig, & Hodapp, 2003). However, results from a meta-analysis of 12 psychophysiological studies found that women had slightly higher heart rates at rest than men and there was a trend of toward higher heart rates during challenge tasks but that men had higher systolic blood pressure in both of these conditions (Stoney, Davis, & Matthews, 1987). In terms of autonomic nervous system responding to acute stress, a recent review by Kajantie and Phillips (2006) concluded that gender differences among adults are marked and consistent; women show lower hypothalamic-pituitary-adrenal axis (HPAA) and autonomic reactivity compared to men.

In sum, evidence for gender differences in physiological reactivity seems to depend on how physiological reactivity is operationalized.

Men show greater responding when reactivity is defined as skin conductance during conditioning/extinction, blood pressure during challenge, and HPA reactivity during challenge; women show greater responding when reactivity is defined as heart rate during rest, and few gender differences are observed when reactivity is operationalized as heart rate during challenge. In addition to methodological differences, another source of variance across these studies is women's hormonal status. Based on their analyses of the age and hormonal status of the sample (when reported) [Kajantie and Phillips \(2006\)](#) concluded that adult women are less reactive than men (lower HPA and autonomic responding) between puberty and menopause, show increased reactivity during the luteal phase and decreased activity during pregnancy and after menopause. Thus, gender differences in physiological reactivity seem to depend on how physiological reactivity is operationalized and may also be influenced by women's hormonal status.

### 2.1.3. Hormonal influences

Several studies examining levels of stress hormones during anxious responding show that men and women exhibit similar cortisol reactivity during acute stress ([Kirschbaum, Kudielka, Gaab, Schommer, & Hellhammer, 1999](#); [Kirschbaum, Wüst, & Hellhammer, 1992](#); [Stoney et al., 1987](#)), although some studies show that men increase their output of adrenaline during acute stress relative to women ([Elwood, Ferguson, & Thakar, 1986](#); [Rauste-von Wright, von Wright, & Frankenhaeuser, 1981](#)).

Hormonal fluctuations related to reproductive events have been advanced as a possible contributor to the gender differences in anxiety. Although [Hedlund and Chambless \(1990\)](#) found no evidence that women are differentially susceptible to conditioning at various times of their menstrual cycle, the hormonal changes related to menstruation and reproductive stages have been associated with increased anxiety among women. For example, state anxiety has been found to increase during the late luteal phase of the menstrual cycle ([Pearlstein et al., 1990](#)) and during pregnancy ([Phillips, Dennerstein, & Farish, 1996](#)). This is consistent with research reviewed by [Kajantie and Philips \(2006\)](#), who noted that during the luteal phase of the menstrual cycle, cortisol output among women is elevated to levels that are similar to men.

According to both retrospective studies ([Williams & Koran, 1997](#)) and prospective studies ([McLeod, Hoehn-Saric, Foster, & Hipsley, 1993](#)), existing anxiety disorders may be exacerbated during the premenstrual phase, as evidenced by increased symptom ratings. Pregnancy among women with panic disorder seems to have a variable effect on symptom intensity. A review of 8 non-controlled studies (all but one were retrospective) by [Hertzberg and Wahlbeck \(1999\)](#) found equivocal results: panic symptoms improved during pregnancy in 41% of women, but that 38% experienced an exacerbation of symptoms. Among women with OCD, a retrospective study by [Williams and Koran \(1997\)](#) found evidence of premenstrual and postpartum exacerbation of symptoms. Pregnancy and postpartum have been identified as periods of increased vulnerability for the development of OCD ([Diaz, Grush, Sichel, & Cohen, 1997](#); [Rasmussen & Eisen, 1990](#); [Ross & McLean, 2006](#)). However, given lack of controlled studies, reliance on retrospective self-report, and the fact that puberty, pregnancy, and childbirth are often periods of considerable stress, it is difficult to attribute disorder onset or exacerbation to co-occurring hormonal changes.

### 2.1.4. Evolutionary influences

In an extensive review of hormonal and physiological data related to the experience of fear and anxiety, [Taylor et al. \(2000\)](#) argued that the familiar *fight-or-flight* characterization of the human stress response does not accurately describe the behavioral response of women. In contrast to the flight-or-flight response, which, as the authors noted, could compromise pregnancy or put women and their

offspring at risk, the authors assert that the female response is marked by a pattern of behavior which they termed *tend-and-befriend*. Tending behaviors include nurturing activities designed to protect the self and offspring, promote safety, and reduce distress, while befriending behaviors create and maintain social networks. According to the evidence reviewed by [Taylor et al. \(2000\)](#), oxytocin and other endogenous opioids appear to underlie these attachment and caregiving behaviors such that when stressed, greater oxytocin production among women has a calming effect that promotes affiliation. Taylor and colleagues support this hypothesis by citing research indicating that higher levels of oxytocin are associated with lower arousal, lower stress responding, and more social behavior. Greater oxytocin production could help to explain findings of weaker adrenocortical responding among male infants ([Davis & Emory, 1995](#)) and adults [Kajantie and Phillips \(2006\)](#) in response to stress cues.

[Craske \(2003\)](#) speculates that need to protect and nurture others, while protective against imminent threat, may help to explain evidence of greater worrying among women when faced with potential threats. A tendency to respond to threat by befriending or by seeking support from others, as opposed to relying on one's own personal coping resources may decrease women's self-efficacy ([Craske, 2003](#)). The role of self-efficacy in the onset and maintenance of anxiety is discussed in greater detail in a later section.

## 2.2. Broad vulnerability factors: higher order factors

### 2.2.1. Negative affectivity

Negative affectivity is strongly linked with both anxiety and mood disorders ([Norton, Sexton, Walker, & Norton, 2005](#)). As expected, negative affectivity is more often observed among girls ([Steiner, Ryst, Berkowitz, Gschwendt, & Koopman, 2002](#)) and adult women across cultures ([Lynn & Martin, 1997](#); [Costa, Terracciano, & McCrae, 2001](#)). During the first few months of life, however, boys show greater negative affect than girls ([Davis & Emory, 1995](#); [Weinberg, Tronick, Cohn, & Olson, 1999](#)). Only at around age two does this pattern reverse, with girls beginning to show a greater propensity to negative affectivity ([Arcus & Kagan, 1995](#)). This shift occurs at approximately the same age in which the effects of gender socialization become evident ([Craske, 2003](#); [Martin, Ruble, & Szkrybalo, 2002](#)). Girls report increasingly higher levels of negative affectivity as they mature, whereas levels remain stable among boys during the same period ([McCrae, Costa, & Terracciano, 2002](#)). Thus, gender differences in negative affectivity become increasingly divergent as the influence of biological factors attenuates relative to the influence of environmental experience ([Lonigan & Phillips, 2001](#)).

### 2.2.2. Trait anxiety

The construct of trait anxiety ([Spielberger, 1966, 1972](#)) is conceptually similar to negative affectivity ([Barlow, 2000](#); [Zinbarg & Barlow, 1996](#)), and women generally score higher on measures of trait anxiety such as the State/Trait Anxiety Inventory (STAI; [Spielberger, Gorsuch, & Lushene, 1970](#)). Among individuals with panic disorder, one study has found greater trait anxiety among women than men with panic disorder ([Chambless & Mason, 1986](#)), but other studies have not replicated this finding ([Turgeon et al., 1998](#); [Foot & Koszycki, 2004](#)) and it has been suggested that the differences detected by [Chambless and Mason \(1986\)](#) may have been statistically but not clinically significant. Possibly, the magnitude of gender differences varies depending on how trait anxiety is measured. For example, [Bander and Betz \(1981\)](#) found few gender differences when using a measure of trait anxiety that emphasizes physiological indicators (the S-R Inventory; [Endler, Hunt, & Rosenstein, 1962](#)), but found that women scored higher than men on the STAI, which reflects cognitive and affective indicators of trait anxiety. A tendency for women to manifest trait anxiety predominately within cognitive/affective domains rather than within the physiological domain is consistent



with studies showing few gender differences in physiological reactivity (Kelly, Forsyth, et al., 2006; Kelly, Tyrka, et al., 2008).

### 2.3. Broad vulnerability factors: disorder specific factors

#### 2.3.1. Anxiety sensitivity

Taylor (1998) and others have emphasized the need for hierarchical models of anxiety to consider disorder-specific risk factors, such as anxiety sensitivity (AS). AS, defined as the fear of anxiety and anxiety-related sensations (Reiss & McNally, 1985), is a risk factor for several anxiety disorders, and is strongly associated with panic disorder (Zinbarg & Barlow, 1996). Several studies have found evidence of greater AS among women within both non-clinical samples of adolescents (Bernstein, Zvolensky, Stewart, Comeau, & Leen-Felder, 2006) and adults (Deacon, Abramowitz, Woods, & Tolin, 2003; Stewart, Taylor, & Baker, 1997; Zvolensky, McNeil, Porter, & Stewart, 2001). Research among clinical samples, however, has been less consistent. Schmidt and Koselka (2000) found significantly greater AS among women with PD than men with PD, whereas Foot and Koszycki (2004) found no gender differences. Measurement of AS, sample characteristics, and methods were similar in both of these studies and therefore, the discrepancy in findings is difficult to interpret.

More consistent gender differences have been found within certain dimensions of AS. Although taxometric studies of AS reveal a similar factor structure for men and women (Dehon, Weems, Stickle, Costa, & Berman, 2005; Stewart et al., 1997), women report greater fear of the physical consequences of anxiety (Stewart et al., 1997), whereas men report greater fear of the social consequences of anxiety (Foot & Koszycki, 2004). The physical concerns factor of AS is considered especially relevant to the etiology of panic (McNally, 1993; Zinbarg, Barlow, & Brown, 1997). Thus, gender differences within a particular dimension of AS may explain an increased risk for panic disorder among women. One possible explanation for this finding is that women objectively experience more physiological symptoms of anxiety than men; women's elevated concern may simply correspond with the extent of their physical symptoms. However, although women report a greater number of symptoms during panic attacks (Dick, Bland, & Newman, 1994; Ströhle, 2000), the mixed findings of research examining gender differences in physiological reactivity do not provide clear support for this hypothesis.

Another possible explanation for gender differences within a particular dimension of AS is that physical symptoms are experienced by men and women equally, but women receive more positive reinforcement for expressing concern toward these symptoms. Over time, such reinforcement could increase self-focused attention that would contribute to actual differences in the experience of physical symptoms. In support of this hypothesis, retrospective studies have found that women are more likely than men to recall being reinforced for expressing somatic concerns (Ehlers, 1993). According to Watt, Stewart, and Cox (1998), men who are discouraged from expressing somatic complaints may suppress them. Instead, they may attend to feared social consequences of visible anxiety symptoms, which could explain why men score higher on the social concerns sub-factor of AS than women (Foot & Koszycki, 2004; Stewart et al., 1997). The mechanisms that account for differences in AS among men and women and how these differences relate to the development of panic disorder have yet to be elucidated.

#### 2.3.2. Disgust sensitivity

Disgust and contamination sensitivity are closely related disorder-specific vulnerability factors that have been associated with animal phobias (e.g., Arrindell, Mulken, & Kok, 1999; Arrindell, 2000). Consistent with the higher prevalence among women for certain types of phobias, several studies indicate that women report greater disgust sensitivity than men (Connolly, Olatunji, & Lohr, 2008; Olatunji, Arrindell, & Lohr, 2005). In fact, Davey (1994) found that the gender

differences in animal fears were largely accounted for by women's higher levels of disgust sensitivity. Similar findings were reported by Olatunji and colleagues (2005), who found that the relationship between gender and blood-injection-injury fears was significantly mediated by differences in disgust sensitivity.

### 2.4. Stress and trauma

#### 2.4.1. Gender-specific trauma

Although women experience fewer traumas than men, they are more likely to experience potentially traumatic events (PTEs) including rape, attempted rape (Koss, Gidycz, & Wisniewski, 1987), and domestic abuse (McHugh, Frieze, & Browne, 1993), that are associated with a high probability of PTSD or other anxiety disorders (Breslau & Anthony, 2007). Women also are more likely than men to experience relationship stressors involving acquaintances, friends, or loved ones, which are referred to as "social network crises" (Kessler & McLeod, 1984). Social network crises may have as much impact on emotional functioning as more commonly identified PTEs (Gold, Marx, Soler-Baillo, & Sloan, 2005). In fact, individuals who have experienced aversive events not subsumed under the DSM-IV-TR definition of a Criterion A traumatic event, such as an unexpected death or serious illness of a loved one, have shown greater PTSD symptomology than individuals who have experienced events classified as traumatic (Gold et al., 2005).

#### 2.4.2. Resiliency

There are several possible explanations for why, even when controlling for trauma type, women are more likely than men to develop PTSD (Kessler et al., 1995; Tolin & Foa, 2006). One explanation is that men and women employ differentially effective coping strategies following exposure to PTEs. Compared to active, problem-focused coping, emotion-based coping strategies (e.g. positive reinterpretation, mental disengagement, denial) are generally less effective in reducing distress (Carver, Scheier, & Weintraub, 1989) and have been associated with greater PTSD symptomology (Blake, Cook, & Keane, 1992). Women tend to use emotion-focused coping more often than men in response to moderately stressful situations (Thoits, 1987). Women's greater tendency to ruminate (discussed further below) may increase their risk for PTSD by inflating the threat-value of PTEs and thereby fostering fearful responding and encouraging avoidance (Davey & Matchett, 1994).

#### 2.4.3. Appraisals of threat

The degree to which men and women experience events as unpredictable, uncontrollable, or overwhelming, may also help to explain why women appear to be more impacted than men by adverse events (Cohen, Kamarck, & Mermelstein, 1983). Compared to men, women are more likely to overestimate the probability of danger (Menzies & Clarke, 1995), to expect harm, and to anticipate poor coping ability (Thorpe & Salkovskis, 1995). From an evolutionary perspective, vigilance toward potential threat and a tendency to judge ambiguous situations as threatening may be hypothesized as adaptive for women by encouraging them to preserve the safety of the self and offspring (Wood & Eagly, 2002). In support of this idea, and parallel to the information processing differences, recent brain imaging studies have found gender differences in the activation of neural structures thought to mediate attention to threat. McClure, Monk, Nelson, Zarahn, and Leibenluft (2004) found that adult women showed greater activation in the orbitofrontal cortex and amygdala when presented with threat cues. No gender differences were observed among adolescents, however, suggesting that the gender differences in patterns of neural responses to emotional faces are not apparent until adulthood.

#### 2.4.4. Uncontrollability

Early childhood experience with diminished control may result in a cognitive style that increases the probability of interpreting events as threatening (Chorpita & Barlow, 1998). Citing research indicating that mother-son dyads demonstrate greater coordination in affective and behavioral displays (e.g., matching facial expressions or direction of attention) than mother-daughter dyads, Craske (2003) suggests that lower synchronicity between mothers and daughters establishes a weaker base of predictability and controllability that puts women at greater risk for anxiety disorders. In adulthood, women report a lower sense of personal control over their lives than men (Ross & Mirowsky, 2002; Thoits, 1987). Thus, in the context of a weaker base of predictability and controllability, women may have fewer opportunities for successful mastery and coping. For example, several studies have shown that caregivers and teachers are more responsive to the actions of boys than girls (Nolen-Hoeksema, 1990). As a result, girls may learn that their behavior has less impact on the environment, which may contribute to a sense of uncontrollability and an attributional style that places them at a greater risk for anxiety and mood disorders (Barlow, 1991; Foa & Riggs, 1994; Nolen-Hoeksema, 1990).

#### 2.4.5. Self-efficacy

Self-efficacy in one's ability to cope with potential threat is a strong predictor of avoidance behavior (Emmelkamp & Felten, 1985). Research among non-clinical samples suggests that women tend to report lower self-efficacy than men (Buchanan & Selmon, 2008; see Bandura, 1997). The manner in which caregivers respond to children's expression of emotion is important to the development of self-efficacy. For example, in a prospective study of 91 mothers of elementary school children who completed a daily checklist of parenting behaviors, Pomerantz and Ruble (1998) found that mothers were more likely to report using control without autonomy-granting with their daughters, but tended to employ control with autonomy-granting with their sons. Furthermore, the greater use of control without autonomy-granting toward girls partly accounted for the finding that girls self-reported a greater tendency take responsibility for failure than boys. The authors suggest that greater caregiver control toward girls may contribute to a heightened vulnerability for anxiety when failure is encountered. Additionally, Krohne and Hock (1991) found that mothers of highly anxious girls tended to maintain control during a problem-solving task, whereas mothers of non-anxious girls were less controlling. This pattern was reversed among boys: mothers of highly anxious boys were less controlling during the problem-solving task than mothers of non-anxious boys. These findings suggest that the influence of caregiver over-control may selectively impact the development and maintenance of anxiety symptoms among girls.

#### 2.4.6. Worry

Worry is a cognitive component of anxiety that is elicited under conditions of uncertainty and which focuses attention toward threat (Borkovec, Shadick, & Hopkins, 1991). Research indicates that most self-reported adult worriers are women (Robichaud, Dugas, & Conway, 2003). With one exception (Suarez & Bell-Dolan, 2001), this pattern of gender difference has also been found among samples of children (Campbell & Rapee, 1994) and preadolescents (Silverman, La Greca, & Wasserstein, 1995). It may be that the incongruent findings relate to methodological differences: Suarez and Bell-Dolan examined worry by children's responses to vignettes, whereas the other two studies involved self-reported worry only. Women's experience with certain uncontrollable life stressors (e.g. greater exposure to particular PTEs) and greater caregiver over-control may support a cognitive style characterized by worry and rumination.

Although worriers do not show deficits in knowledge of problem-solving skills, they tend to have a negative problem orientation,

defined as low problem-solving confidence and low perceived control over the problem-solving process in the absence of actual problem-solving skills deficits. Research among non-clinical samples indicates that women's greater tendency to worry may be due, in part, to a more negative problem orientation (Robichaud et al., 2003). Women in this study reported greater worry and a more negative problem orientation than men. However, the gender difference in worry disappeared when controlling for women's more negative problem orientation. Thus, women's approach to problem-solving appears to be an important avenue for understanding how worry contributes to anxiety among women.

#### 2.4.7. Rumination

A similar construct to worry, rumination is defined as thoughts or behaviors that focus attention toward depressive symptoms or their consequences, rather than toward pleasant or neutral topics (Nolen-Hoeksema, 1991). Rumination has been found to prolong anxious moods (Blagden & Craske, 1996). Compared to men, women are more likely to ruminate in response to dysphoric mood (Nolen-Hoeksema, 1991). Further, greater rumination among women is not attributable to higher levels of distress, greater emotional expressivity, or greater willingness to endorse socially undesirable traits (Nolen-Hoeksema & Jackson, 2001). Instead, gender differences in rumination are mediated by three beliefs, all of which were endorsed more by women than by men: negative emotions are difficult to control, a positive emotional tone of interpersonal relationships is a personal responsibility, and negative events are difficult to control (Nolen-Hoeksema & Jackson, 2001). It is not known whether women ruminate more than men in response to anxious mood. However, their tendency to focus attention on negative interpretations and outcome expectancies would likely increase state anxiety.

#### 2.4.8. Heightened sensitivity to social cues

Compared to men, women are generally better at identifying affect in nonverbal cues (Hall, 1987). For example, gender differences in reactivity to facial expressions emerge at the same developmental time as differences in negative affectivity and socialization appear (Craske, 2003). Because facial expressions are a primary means of communicating danger, women's greater sensitivity to facial expressions places them at particular risk for developing fears and phobias (Craske, 2003). A vulnerability to socially-transmitted threat information may be especially detrimental to girls reared by anxious caregivers, given that they have more opportunities to learn about threat and avoidance. In fact, a study by Chorpita, Albano, and Barlow (1996) found that family discussions of ambiguously threatening situations led to increased avoidance behavior among anxious children (differences between boys and girls were not examined). Greater fear and anxiety among women may be influenced by two related factors. First, women may be more prone to vicarious fear conditioning because they are more sensitive to socially-transmitted information and are more attentive to others' expressions of fear. Second, women's distress following aversive experiences may be inflated subsequent to ruminative reevaluation of the threat-value of stimuli.

#### 2.5. Behavioral avoidance

Avoidance is central to onset and maintenance of anxiety disorders (Barlow, 2001; Mowrer, 1960). Gender socialization that encourages exposure to feared stimuli facilitates the emotional processing of feared stimuli. Gender differences in fears and phobias increase throughout childhood, which may reflect differential extinction rates to developmentally appropriate fears (Fredrikson, Annas, & Fischer, 1996). If boys are more encouraged to confront fears and more dissuaded to avoid feared situations than girls, this reinforcement may motivate behavior that affords opportunities for emotional processing of fears. In this context, evidence showing that there are few gender

differences in social fears relative to other types of anxiety may be due to equal levels of reinforcement for approaching social situations. Reinforcement patterns that support avoidance among girls may thwart opportunities for them to emotionally process fears, thereby preventing extinction of existing fears and inhibiting the development of self-efficacy.

## 2.6. Environmental influences

### 2.6.1. Sociocultural influences

Coping strategies are shaped by social, cultural, and developmental contexts that influence men and women differently. As noted by [Chambless \(1989\)](#), in Western societies, the typical age of onset for agoraphobia corresponds with a developmental period marked by separation issues (e.g., leaving the home, beginning a career) and the adoption of multiple roles and obligations, which may be more difficult for women. Difficulty coping with separation, especially when combined with low autonomy, is a risk factor for agoraphobia ([Chambless & Goldstein, 1981](#)), which suggests that agoraphobia may be less common in cultures where separation issues are minimized and role obligations are more limited. In some non-Western cultures, in which it is more culturally acceptable for women to stay home (unless accompanied by an adult male), symptoms of agoraphobia are reportedly low ([Kirmayer, Young, & Hayton, 1995](#)). For example, [El-Islam \(1994\)](#) found that women in Qatar reported lower levels of agoraphobia than women in the U.S. The authors hypothesized that this cross-cultural discrepancy may be due to the fewer, more clearly defined social role obligations in the Qatari culture than in American culture.

[Arrindell and colleagues \(2003\)](#) examined how agoraphobic fears relate to masculinity measured at a national level across 11 countries. National masculinity was defined as the degree to which the society delineates distinct and rigid gender roles and upholds strong patriarchal values. A significant relationship was found between national levels of masculinity and agoraphobic fears, such that the greater the rigidity in gender roles at the sociocultural level, the more likely that men and women endorsed agoraphobic fears. Patriarchal societies that de-emphasize assertiveness and independence among women may create sociocultural contexts that foster fearfulness and avoidant coping ([Arrindell et al., 2003](#)). This conclusion appears to be at odds with the findings of [El-Islam \(1994\)](#) who found that low agoraphobic fears were related to the limited, defined social role obligations of patriarchal Qatar. It could be that agoraphobic fears were low among the Qatari women because the women in this culture have few opportunities to leave their home unaccompanied. However, given that [Arrindell and colleagues \(2003\)](#) did not sample women from any Middle Eastern countries and that there is evidence of considerable variation in gender roles across cultures ([Basow, 1984](#); [Chia, Moore, Lam, Chuang, & Cheng, 1994](#)), it would be premature to draw conclusions regarding these discrepant findings.

### 2.6.2. Gender role socialization

Social patterns of reinforcement for girls and boys differ in ways that are critical to understanding the gender differences in anxiety. [Bem's \(1981\)](#) gender role theory asserts that boys and girls are socialized to develop socially prescribed behaviors, traits, skills, and interests that are consistent with their gender. Thus, because the expression of anxiety is inconsistent with the male gender role, fearful behavior may be less tolerated in boys. Caregivers and other socialization agents (e.g., teachers, peers, and media) may encourage gender conforming behaviors by differentially reinforcing agency and assertiveness among boys and anxious behaviors among girls. Research indicates that parents consider withdrawal and inhibition to be less acceptable as boys get older, but not for girls ([Stevenson-Hinde & Shouldice, 1993](#)).

There is considerable evidence demonstrating that gender role is significantly related to symptoms of anxiety and fear ([Chambless &](#)

[Mason, 1986](#); [Ollendick, King, & Muris, 2002](#)). Research consistently has found a significant association between scores on standard fear scales, such as the Fear Survey Schedule ([Wolpe & Lang, 1977](#)) or the Fear Survey Schedule for Children-II ([Gullone & King, 1993](#)), and gender role scales, such as the Bem Sex Role Inventory, which measure the degree to which individuals endorse personality characteristics consistent with social conceptions of masculine and feminine gender roles. Among children, greater fear reporting has been associated with higher levels of femininity ([Muris, Meesters, & Knoop, 2005](#)) and lower levels of masculinity ([Ginsburg & Silverman, 2000](#)). Similar findings are seen in adults, with some studies demonstrating a positive relationship between fear and femininity ([Dillon, Wolf, & Katz, 1985](#); [Tucker & Bond, 1997](#)), others showing a negative association with masculinity ([Arrindell, 2000](#)), and others finding that both high femininity and low masculinity are related to elevated fear ([Carey, Dusek, & Spector, 1988](#)). Several studies have found a significant relationship between femininity and trait anxiety ([Bander & Betz, 1981](#); [Biaggio & Nielsen, 1976](#); [Gall, 1969](#)).

Additionally, [Bander and Betz \(1981\)](#) found that gender differences were greater for math anxiety (a gender-stereotypic domain) than in test anxiety (a gender-neutral domain), suggesting that the magnitude of gender difference depends on the type of anxiety. [Todman and Day \(2006\)](#) found that gender role significantly predicted computer-related anxiety even when controlling for gender and department type (arts vs. engineering), revealing that gender differences in certain domain-specific types of anxiety may relate to gender role socialization.

[Moscovitch, Hofmann, and Litz \(2005\)](#) found that masculinity was significantly negatively related to social anxiety, whereas biological sex was unrelated to social anxiety. This study also found that self-perceived levels of independence and interdependence predicted social anxiety differently among men and women. In men, lower levels of independence and higher levels of interdependence were associated with greater social anxiety, whereas this pattern was reversed among women. These findings may support a self-discrepancy theory in which social anxiety increases among men and women when self-perceptions run counter to social mores. Thus, individuals who can be flexible across situations, regardless of the gender-related social demands of the situation, feel less social anxiety and discomfort. This is consistent with data showing that androgynous women (who are likely to be flexible across situations) report the lowest levels of social-evaluative fears ([Goodman & Kantor, 1983](#)). Further examination of how perceived gender role and ideal gender role relate with social anxiety may help explain why gender differences in social anxiety are markedly small, relative to other anxiety disorders.

If expression of fear and avoidance is less acceptable among men, boys may learn that the masculine role involves bravery and purposeful coping behavior when faced with anxiety-provoking situations ([Bem, 1981](#)). [Greif, Alvarez, and Ullman \(1981\)](#) found that mothers are more likely to discuss emotional states with their daughters, whereas they are more likely to discuss the causes and consequences of the feelings with their sons. Boys are thus encouraged to focus on problem-solving and gaining control over their emotion, rather than on the experience of the emotion itself. Learning to cope with anxiety in this problem-focused manner may help equip men with the instrumental traits and skills that prevent excessive fears or other anxiety disorders from developing. In contrast, a traditional feminine gender role that deemphasizes autonomy and mastery while promoting dependency and expectations of protection would be more compatible with avoidance behavior.

## 3. Conclusion

Substantial evidence points to a preponderance of women demonstrating greater fear and anxiety than men across the life span. Many of



the risk factors for developing anxiety form the basis for the expected gender differences. For example, women tend to score higher on higher-order vulnerability factors including negative affectivity and trait anxiety. Further, genetic research suggests that these factors are more heritable among women for reasons that are still unclear. One possibility is that environmental factors counteract anxious tendencies among boys but support them among girls. Although few temperamental differences are observed in very young infants, there is a shift toward greater negative affectivity among girls during the developmental period at which gender socialization becomes evident. Timing of this shift implicates environmental influences such as gender socialization.

Much of the hormonal and physiological data for gender differences appear contradictory. Several studies suggest that men are somewhat more physiologically reactive than women in response to acute stressors. While seemingly inconsistent with broad gender differences in anxiety, this finding is consistent with women's greater tendency to worry, which is known to decrease physiological reactivity. It is also consistent with evidence that oxytocin production among women leads to lower arousal and stress responding. Evidence that men are more responsive to acute threat, whereas women are more distressed by potential threat also fits well with the tend-and-befriend stress response articulated by Taylor et al. (2000). However, through hypervigilance to threat and prioritizing social networks, the tend-and-befriend response may lower women's perceived coping ability and encourage avoidance.

Although women are not more prone to experience traumas overall, they are more likely to experience certain types of trauma, including sexual abuse and social network crises, which may be particularly potent risk factors for anxiety. Further, in response to aversive events, women are more likely to use emotion-focused coping which is less effective in reducing distress than problem-focused coping strategies.

Through lower parent-child synchronicity and caregiver overprotection, girls may have less opportunity to exert control over their environment. As a result, women may perceive themselves as less capable of coping with potentially threatening situations. Lower self-efficacy may, in turn, motivate women to rely on a threat-based style of responding and to cope with uncertainty through avoidance. Avoidant coping among women is consistent with traditional gender roles, in which men and women are expected to respond to potential threat differently. Thus, the female stress response, as viewed from an evolutionary perspective that emphasizes the need to protect and nurture offspring, may be reinforced by gender socialization processes that support worry, sensitivity to socially-transmitted information, and avoidant coping.

In sum, gender differences in fear and anxiety are greatest when 1) examining molar levels versus biological levels of analysis and 2) examining individuals after the age at which socialization factors begin to exert influence. Genetic vulnerabilities gradually evolve into fully articulated traits through complex, bidirectional interactions with environmental factors. Thus, gender differences at each level of analysis are likely moderated by socialization processes that prescribe gender-specific expectations regarding the expression of anxiety and the acceptable means of coping with anxiety. These socialization factors influence expression of traits by shaping patterns of reinforcement that cultivate and promote processes related to anxiety.

## References

- Anderson, J. C., Williams, S. M., McGee, R., & Silva, P. A. (1987). DSM-III disorders in preadolescent children: Prevalence in a large sample from the general population. *Archives of General Psychiatry*, 44(1), 69–76.
- Angst, J., & Dobler-Mikola, A. (1985). The Zurich Study: V. Anxiety and phobia in young adults. *European Archives of Psychiatry and Neurological Sciences*, 235(3), 171–178.
- Arcus, D., & Kagan, J. (1995). Temperament and craniofacial variation in the first two years. *Child Development*, 66(5), 1529–1540.
- Arrindell, W. A. (2000). Phobic dimensions: IV. The structure of animal fears. *Behaviour Research and Therapy*, 38(5), 509–530.
- Arrindell, W. A., Eisemann, M., Richter, J., Oei, T. P. S., Caballo, V. E., Ende, J., et al. (2003). Masculinity-femininity as a national characteristic and its relationship with national agoraphobic fear levels: Fodor's sex role hypothesis revitalized. *Behaviour Research and Therapy*, 41, 795–807.
- Arrindell, W. A., Mulken, S., & Kok, J. (1999). Disgust sensitivity and the sex differences in fears to common indigenous animals. *Behaviour Research and Therapy*, 37(3), 273–280.
- Basow, S. A. (1984). Cultural variations in sex-typing. *Sex Roles*, 10, 577–585.
- Bander, R. S., & Betz, N. E. (1981). The relationship of sex and sex role to trait and situationally specific anxiety types. *Journal of Research in Personality*, 15(3), 312–322.
- Bandura, A. (1997). *Self-efficacy: The exercise of control*. New York: Freeman.
- Barlow, D. H. (1991). Disorders of emotion. *Psychological Inquiry*, 2(1), 58–71.
- Barlow, D. H. (2000). Unraveling the mysteries of anxiety and its disorders from the perspective of emotion theory. *American Psychologist*, 55(11), 1247–1263.
- Barlow, D. H. (2001). *Anxiety & its disorders*, 2nd ed. New York: Guilford.
- Bem, S. L. (1981). Gender schema theory: A cognitive account of sex typing. *Psychological Review*, 88(4), 354–364.
- Bernstein, A., Zvolensky, M. J., Stewart, S. H., Comeau, M. N., & Leen-Felder, E. W. (2006). Anxiety sensitivity taxonomic across gender among youth. *Behaviour Research and Therapy*, 44(5), 679–698.
- Biaggio, M. K., & Nielsen, E. C. (1976). Anxiety correlates of sex-role identity. *Journal of Clinical Psychology*, 32(3), 619–623.
- Blagden, J. C., & Craske, M. G. (1996). Effects of active and passive rumination and distraction: A pilot replication with anxious mood. *Journal of Anxiety Disorders*, 10, 243–252.
- Blake, D. D., Cook, J. D., & Keane, T. M. (1992). Post-traumatic stress disorder and coping in veterans who are seeking medical treatment. *Journal of Clinical Psychology*, 48, 695–704.
- Borkovec, T., Shadick, R., & Hopkins, M. (1991). *The nature of normal and pathological worry. Chronic anxiety: Generalized anxiety disorder and mixed anxiety-depression* (pp. 29–51). New York, NY US: Guilford Press.
- Bourdon, K. H., Boyd, J. H., Rae, D. S., Burns, B. J., Thompson, J. W., & Lock, B. Z. (1988). Gender differences in phobias: Results of the ECA community survey. *Journal of Anxiety Disorders*, 2(3), 227–241.
- Bowen, R. C., Offord, D. R., & Boyle, M. H. (1990). The prevalence of overanxious disorder and separation anxiety disorder: Results from the Ontario Child Health Study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 29(5), 753–758.
- Breslau, N., & Anthony, J. C. (2007). Gender differences in the sensitivity to posttraumatic stress disorder: An epidemiological study of urban young adults. *Journal of Abnormal Psychology*, 116, 607–611.
- Breslau, N., Chilcoat, H. D., Peterson, E. L., & Schultz, L. R. (2000). Gender differences in major depression: The role of anxiety. In E. Frank (Ed.), *Gender and its effects on psychopathology* (pp. 131–150). Washington, DC, US: American Psychiatric Publishing, Inc.
- Bruce, S. E., Yonkers, K. A., Otto, M. W., Eisen, J. L., Weisberg, R. B., Pagano, M., et al. (2005). Influence of psychiatric comorbidity on recovery and recurrence in generalized anxiety disorder, social phobia, and panic disorder: A 12-year prospective study. *American Journal of Psychiatry*, 162(6), 1179–1187.
- Buchanan, T., & Selmon, N. (2008). Race and gender differences in self-efficacy: Assessing the role of gender role attitudes and family background. *Sex Roles*, 58(11–12), 822–836.
- Cameron, O. G., & Hill, E. M. (1989). Women and anxiety. *Psychiatric Clinics of North America*, 12(1), 175–186.
- Campbell, M. A., & Rapee, R. M. (1994). The nature of feared outcome representations in children. *Journal of Abnormal Child Psychology*, 22(1), 99–111.
- Carey, M. P., Dusek, J. B., & Spector, I. P. (1988). Sex roles, gender and fears: A brief report. *Phobia Practice and Research Journal*, 1(1), 114–120.
- Carver, C. S., Scheier, M. F., & Weintraub, J. K. (1989). Assessing coping strategies: A theoretically based approach. *Journal of Personality and Social Psychology*, 56, 267–283.
- Castle, D. J., Deale, A., & Marks, I. M. (1995). Gender differences in obsessive compulsive disorder. *Australian and New Zealand Journal of Psychiatry*, 29(1), 114–117.
- Chambless, D. L. (1989). Agoraphobia. In C. Lindemann & M. D. Latham (Eds.), *Handbook of phobia therapy: Rapid symptom relief in anxiety disorders* Northvale, NJ: Jason Aronson, Inc.
- Chambless, D. L., & Goldstein, A. J. (1981). Clinical treatment of agoraphobia. In M. Mavissakalian & D. H. Barlow (Eds.), *Phobia: Psychological and pharmacological treatment* (pp. 103–144). New York: Guilford.
- Chambless, D. L., & Mason, J. (1986). Sex, sex-role stereotyping and agoraphobia. *Behaviour Research and Therapy*, 24(2), 231–235.
- Chia, R. C., Moore, I. L., Lam, K. N., Chuang, C. J., & Cheng, B. S. (1994). Cultural differences in gender role attitudes between Chinese and American students. *Sex Roles*, 31, 23–30.
- Chorpita, B. F., Albano, A. M., & Barlow, D. H. (1996). Cognitive processing in children: Relation to anxiety and family influences. *Journal of Clinical Child Psychology*, 25, 170–176.
- Chorpita, B. F., & Barlow, D. H. (1998). The development of anxiety: The role of control in the early environment. *Psychological Bulletin*, 124, 3–21.
- Cohen, S., Kamarck, T., & Mermelstein, R. A. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24, 385–396.
- Connolly, K. M., Olatunji, B. O., & Lohr, J. M. (2008). Evidence for disgust sensitivity mediating the sex differences found in blood-injection-injury phobia and spider phobia. *Personality and Individual Differences*, 44, 898–908.
- Costa, P., Jr., Terracciano, A., & McCrae, R. R. (2001). Gender differences in personality traits across cultures: Robust and surprising findings. *Journal of Personality and Social Psychology*, 81(2), 322–331.



- Costello, E. J., Egger, H. L., & Angold, A. (2003). Development epidemiology of anxiety disorders. In T. H. Ollendick & J. S. March (Eds.), *Phobic Anxiety Disorders in Children and Adolescents: A Clinician's Guide to Effective Psychosocial and Pharmacological Interventions* (pp. 61–91). Oxford, UK: Oxford University Press.
- Cox, B. J., Swinson, R. P., & Shulman, I. D. (1993). Gender effects and alcohol use in panic disorder with agoraphobia. *Behaviour Research and Therapy*, 31(4), 413–416.
- Craig, K. D., & Lowery, H. J. (1969). Heart-rate components of conditioned vicarious autonomic responses. *Journal of Personality and Social Psychology*, 11(4), 381–387.
- Craske, M. G. (2003). The origins of phobias and anxiety disorders: Why more women than men. Oxford, England: Elsevier Science.
- Davey, G. C. L. (1994). Self-reported fears to common indigenous animals in an adult UK population: The role of disgust sensitivity. *British Journal of Psychology*, 85(4), 541–554.
- Davey, G. C. L., & Matchett, G. (1994). Unconditioned stimulus rehearsal and the retention and enhancement of differential "fear" conditioning: Effects of trait and state anxiety. *Journal of Abnormal Psychology*, 103, 708–718.
- Davey, G. C. L., McDonald, A. S., & Hirisave, U. (1998). A cross-cultural study of animal fears. *Behaviour Research and Therapy*, 36(7–8), 735–750.
- Davis, M., & Emory, E. (1995). Sex differences in neonatal stress reactivity. *Child Development*, 66(1), 14–27.
- Deacon, B. J., Abramowitz, J. S., Woods, C. M., & Tolin, D. F. (2003). The anxiety sensitivity index – Revised: Psychometric properties and factor structure in two nonclinical samples. *Behaviour Research and Therapy*, 41(12), 1427–1449.
- Dehon, C., Weems, C. F., Stickle, T. R., Costa, N. M., & Berman, S. L. (2005). A cross sectional evaluation of factorial invariance of anxiety sensitivity in adolescents and young adults. *Behavior Research and Therapy*, 43(6), 799–810.
- Diaz, S. F., Grush, L. R., Sichel, D. A., & Cohen, L. S. (1997). Obsessive-compulsive disorder in pregnancy and the puerperium. In L. J. Dickstein, M. B. Riba, & J. M. Oldham (Eds.), *American Psychiatric Press review of psychiatry*, vol. 16. (pp. III–97–III–112) Washington, DC, US: American Psychiatric Association.
- Dick, C. L., Bland, R. C., & Newman, S. C. (1994). Panic disorder. *Acta Psychiatrica Scandinavica*, 89(376, Suppl.), 45–53.
- Dillon, K. M., Wolf, E., & Katz, H. (1985). Sex roles, gender, and fear. *Journal of Psychology*, 119, 355–359.
- Ehlers, A. (1993). Somatic symptoms and panic attacks: A retrospective study of learning experiences. *Behaviour Research and Therapy*, 31(3), 269–278.
- Eley, T. C. (2001). Contributions of behavioral genetics research: Quantifying genetic, shared environmental and nonshared environmental influences. In M. D. Vasey & R. Mark (Eds.), *The developmental psychopathology of anxiety* (pp. 45–59). New York, NY, US: Oxford University Press.
- El-Islam, M. F. (1994). Cultural aspects of morbid fears in Qatari women. *Social Psychiatry and Psychiatric Epidemiology*, 29(3), 137–140.
- Elwood, S. W., Ferguson, H. B., & Thakar, J. (1986). Catecholamine response of children in a naturally occurring stressor situation. *Journal of Human Stress*, 12(4), 154–161.
- Emmelkamp, P. M. G., & Felten, M. (1985). The process of exposure in vivo: cognitive and physiological changes during treatment of acrophobia. *Behaviour Research and Therapy*, 23, 219–223.
- Endler, N. S., Hunt, J. M., & Rosenstein, A. J. (1962). An S–R inventory of anxiousness. *Psychological Monographs*, 76(17, Whole No. 536), 33.
- Foa, E. B., & Riggs, D. S. (1994). Posttraumatic stress disorder and rape. In R. S. Pynoos (Ed.), *Posttraumatic stress disorder: A clinical review* Baltimore, MD, US: The Sidran Press.
- Foot, M., & Koszycki, D. (2004). Gender differences in anxiety-related traits in patients with panic disorder. *Depression and Anxiety*, 20(3), 123–130.
- Fredrikson, M., Annas, P., & Fischer, H. (1996). Gender and age differences in the prevalence of specific fears and phobias. *Behaviour Research and Therapy*, 34(1), 33–39.
- Gall, M. D. (1969). The relationship between masculinity-femininity and manifest anxiety. *Journal of Clinical Psychology*, 25(3), 294–295.
- Gater, R., Tansella, M., Korten, A., Tiemens, B. G., Mavreas, V. G., & Olatawura, M. O. (1998). Sex differences in the prevalence and detection of depressive and anxiety disorders in general health care settings: Report from the World Health Organization collaborative study on psychological problems in general health care. *Archives of General Psychiatry*, 55(5), 405–413.
- Ginsburg, G. S., & Silverman, W. K. (2000). Gender role orientation and fearfulness in children with anxiety disorders. *Journal of Anxiety Disorders*, 14(1), 57–67.
- Gold, S. D., Marx, B. P., Soler-Baillo, J. M., & Sloan, D. M. (2005). Is life stress more traumatic than traumatic stress? *Journal of Anxiety Disorders*, 19(6), 687–698.
- Goldstein, R. B., Wickramaratne, P. J., Horwath, E., & Weissman, M. M. (1997). Familial aggregation and phenomenology of 'early-onset (at or before age 20 years) panic disorder. *Archives of General Psychiatry*, 54(3), 271–278.
- Goodman, S., & Kantor, D. (1983). Influence of sex-role identity on two indices of social anxiety. *Journal of Research in Personality*, 17, 443–450.
- Graham, L. A., Cohen, S. I., & Shmavonian, R. M. (1966). Sex differences in autonomic responses during instrumental conditioning. *Psychosomatic Medicine*, 28(3), 264–271.
- Greif, E., Alvarez, M., & Ulman, K. (1981). Recognizing emotions in other people: Sex differences in socialization. *Poster presented at the biennial meeting of the Society for Research in Child Development*, Boston, MA.
- Gullone, E., & King, N. J. (1993). The fears of youth in the 1990s: Contemporary normative data. *Journal of Genetic Psychology*, 154, 137–153.
- Hall, J. A. (1987). On explaining gender differences: The case of nonverbal communication. In P. Shaver & C. Hendrick (Eds.), *Sex and gender* Thousand Oaks, CA, US: Sage Publications, Inc.
- Hedlund, M. A., & Chambless, D. L. (1990). Sex differences and menstrual cycle effects in aversive conditioning: A comparison of premenstrual and intermenstrual women with men. *Journal of Anxiety Disorders*, 4(3), 221–231.
- Hertzberg, T., & Wahlbeck, K. (1999). The impact of pregnancy and puerperium on panic disorder: A review. *Journal of Psychosomatic Obstetrics and Gynecology*, 20, 59–64.
- Hettema, J. M., Prescott, C. A., Myers, J. M., Neale, M. C., & Kendler, K. S. (2005). The structure of genetic and environmental risk factors for anxiety disorders in men and women. *Archives of General Psychiatry*, 62(2), 182–189.
- Himle, J. A., McPhee, K., Cameron, O. G., & Curtis, G. C. (1989). Simple phobia: Evidence for heterogeneity. *Psychiatry Research*, 28(1), 25–30.
- Hollifield, M., Katon, W., Skipper, B., Chapman, T., Ballenger, J. C., Mannuzza, S., et al. (1997). Panic disorder and quality of life: Variables predictive of functional impairment. *American Journal of Psychiatry*, 154(6), 766–772.
- Kajantie, E., & Phillips, D. I. (2006). The effects of sex and hormonal status on the physiological response to acute psychosocial stress. *Psychoneuroendocrinology*, 31, 151–178.
- Karno, M., & Golding, J. M. (1991). Obsessive compulsive disorder. In L. N. Robins & D. A. Regier (Eds.), *Psychiatric disorders in America* (pp. 204–219). New York: Free Press.
- Karno, M., Golding, J. M., Sorenson, S. B., & Burman, M. A. (1988). The epidemiology of obsessive-compulsive disorder in five US communities. *Archives of General Psychiatry*, 45(12), 1094–1099.
- Katkin, E. S., & Hoffman, L. S. (1976). Sex differences and self-report of fear: A psychophysiological assessment. *Journal of Abnormal Psychology*, 85(6), 607–610.
- Katzelnick, D. J., & Greist, J. H. (2001). Social anxiety disorder: An unrecognized problem in primary care. *Journal of Clinical Psychiatry*, 62(Suppl1), 11–16.
- Kelly, M. M., Forsyth, J. P., & Karekla, M. (2006). Sex differences in response to a panicogenic biological challenge procedure: An experimental evaluation of panic vulnerability in a non-clinical sample. *Behaviour Research and Therapy*, 44, 1421–1430.
- Kelly, M. M., Tyrka, A. R., Anderson, G. M., Price, L. H., & Carpenter, L. L. (2008). Sex differences in emotional and physiological responses to the Trier Social Stress Test. *Journal of Behavior Therapy and Experimental Psychiatry*, 39(1), 87–98.
- Kendler, K. S., Neale, M. C., Kessler, R. C., Heath, A. C., & Eaves, L. J. (1992). The genetic epidemiology of phobias in women: The interrelationship of agoraphobia, social phobia, situational phobia, and simple phobia. *Archives of General Psychiatry*, 49(4), 273–281.
- Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C., Hughes, M., Eshelman, S., et al. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Study. *Archives of General Psychiatry*, 51(1), 8–19.
- Kessler, R. C., & McLeod, J. D. (1984). Sex differences in vulnerability to undesirable life events. *American Sociological Review*, 49(5), 620–631.
- Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, 52(12), 1048–1060.
- Kirschbaum, C., Kudielka, B. M., Gaab, J., Schommer, N. C., & Hellhammer, D. H. (1999). Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosomatic Medicine*, 61, 154–162.
- Kirschbaum, C., Wust, S., & Hellhammer, D. (1992). Consistent sex differences in cortisol responses to psychological stress. *Psychosomatic Medicine*, 54, 648–657.
- Kirmayer, L. J., Young, A., & Hayton, B. C. (1995). The cultural context of anxiety disorders. *Psychiatric Clinics of North America*, 18(3), 503–521.
- Klorman, R. (1974). Psychometric description of some specific-fear questionnaires. *Behavior Therapy*, 5(3), 401–409.
- Koss, M. P., Gidycz, C. A., & Wisniewski, N. (1987). The scope of rape: Incidence and prevalence of sexual aggression and victimization in a national sample of higher education students. *Journal of Consulting and Clinical Psychology*, 55(2), 162–170.
- Krohne, H. W., & Hock, M. (1991). Relationships between restrictive mother-child interactions and anxiety of the child. *Anxiety Research*, 4(2), 109–124.
- Lake, R. I. E., Eaves, L. J., Maes, H. H. M., Heath, A. C., & Martin, N. G. (2000). Further evidence against the environmental transmission of individual differences in neuroticism from a collaborative study of 45,850 twins and relatives on two continents. *Behavior Genetics*, 30(3), 223–233.
- Lewinsohn, P. M., Gotlib, I. H., Lewinsohn, M., Seeley, J. R., & Allen, N. B. (1998). Gender differences in anxiety disorders and anxiety symptoms in adolescents. *Journal of Abnormal Psychology*, 107, 109–117.
- Líndal, E., & Stefánsson, J. G. (1993). The lifetime prevalence of anxiety disorders in Iceland as estimated by the US National Institute of Mental Health Diagnostic Interview Schedule. *Acta Psychiatrica Scandinavica*, 88(1), 29–34.
- Lonigan, C. J., & Phillips, B. M. (2001). Temperamental basis of anxiety disorders in children. In M. W. Vasey & M. R. Dadds (Eds.), *The developmental psychopathology of anxiety* (pp. 60–91). New York: Oxford University Press.
- Lynn, R., & Martin, T. (1997). Gender differences in extraversion, neuroticism, and psychoticism in 37 nations. *Journal of Social Psychology*, 137(3), 369–373.
- Maltzman, I. (1979, June). Habituation of the GSR and digital vasomotor components of the orienting reflex as a consequence of task instructions and sex differences. *Physiological Psychology*, 7(2), 213–220.
- Martin, C. L., Ruble, D. N., & Szkrybalo, J. (2002). Cognitive theories of early gender development. *Psychological Bulletin*, 128(6), 903–933.
- McClure, E. B., Monk, C. S., Nelson, E. E., Zarahn, E., Leibenluft, E., Bilder, R. M., et al. (2004). A developmental examination of gender differences in brain engagement during evaluation of threat. *Biological Psychiatry*, 55(11), 1047–1055.
- McCrae, R. R., Costa, P. T., Jr., & Terracciano, A. (2002). Personality trait development from age 12 to age 18: Longitudinal, cross-sectional and cross-cultural analyses. *Journal of Personality and Social Psychology*, 83(6), 1456–1468.
- McGee, R., Feehan, M., Williams, S., Partridge, F., Silva, P. A., & Kelly, J. (1990). DSM-III disorders in a large sample of adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 29(4), 611–619.
- McHugh, M. C., Frieze, I. H., & Browne, A. (1993). Research on battered women and their assailants. In F. L. Denmark & M. A. Paludi (Eds.), *Psychology of women: A handbook*

- of issues and theories. CT Westport US: Greenwood Press/Greenwood Publishing Group, Inc.
- McLeod, D. R., Hoehn-Saric, R., Foster, G. V., & Hipsley, P. A. (1993). The influence of premenstrual syndrome on ratings of anxiety in women with generalized anxiety disorder. *Acta Psychiatrica Scandinavica*, 88, 248–251.
- McNally, R. J. (1993). *Panic disorder: A critical analysis*. New York, NY, US: Guilford Press.
- Menzies, R. G., & Clarke, J. C. (1995). The etiology of phobias: A non-associative account. *Clinical Psychology Review*, 15, 23–48.
- Mowrer, O. H. (1960). *Learning theory and behavior*. New York: Wiley.
- Moscovitch, D. A., Hofmann, S. G., & Litz, B. T. (2005). The impact of self-construals on social anxiety: A gender-specific interaction. *Personality and Individual Differences*, 38, 659–672.
- Muris, P., & Merckelbach, H. (2001). The etiology of childhood specific phobias: A multifactorial model. In M. W. Vasey & M. Dadds (Eds.), *The developmental psychopathology of anxiety* Oxford University Press: New York.
- Muris, P., Meesters, C., & Knoop, M. (2005). The relation between gender role orientation and fear and anxiety in nonclinic-referred children. *Journal of Clinical Child and Adolescent Psychology*, 34, 326–332.
- Muris, P., & Ollendick, T. H. (2002). The assessment of contemporary fears in adolescents using a modified version of the Fear Survey Schedule for Children-Revised. *Journal of Anxiety Disorders*, 16(6), 567–584.
- Nolen-Hoeksema, S. (1990). *Sex Differences in Depression*. Stanford, CA: Stanford University Press.
- Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology*, 100(4), 569–582.
- Nolen-Hoeksema, S., & Jackson, B. (2001). Mediators of the gender difference in rumination. *Psychology of Women Quarterly*, 25(1), 37–47.
- Norton, P. J., Sexton, K. A., Walker, J. R., & Norton, R. G. (2005). Hierarchical model of vulnerabilities for anxiety: Replication and extension with a clinical sample. *Cognitive Behaviour Therapy*, 34(1), 50–63.
- Ochoa, L., Beck, A. T., & Steer, R. A. (1992). Gender differences in comorbid anxiety and mood disorders. *American Journal of Psychiatry*, 149(10), 1409–1410.
- Olatunji, B. O., Arrindell, W. A., & Lohr, J. M. (2005). Can the sex differences in disgust sensitivity account for the sex differences in blood-injection-injury fears? *Personality and Individual Differences*, 39(1), 61–71.
- Ollendick, T. H. (1983). Reliability and validity of the Revised Fear Survey Schedule for Children (FSSC-R). *Behaviour Research and Therapy*, 21(6), 685–692.
- Ollendick, T. H., King, N. J., & Muris, P. (2002). Fears and phobias in children: Phenomenology, epidemiology, and aetiology. *Child & Adolescent Mental Health*, 7, 98–106.
- Pearlstein, T. B., Frank, E., Rivera-Tovar, A., Thoft, J. S., Jacobs, E., & Mieczkowski, T. A. (1990). Prevalence of Axis I and Axis II disorders in women with late luteal phase dysphoric disorder. *Journal of Affective Disorders*, 20(2), 129–134.
- Perna, G., Caldirola, D., Arancio, C., & Bellodi, L. (1997). Panic attacks: A twin study. *Psychiatry Research*, 66(1), 69–71.
- Perna, G., Bertani, A., Caldirola, D., & Bellodi, L. (1996). Family history of panic disorder and hypersensitivity to CO-sub-2 in patients with panic disorder. *American Journal of Psychiatry*, 153(8), 1060–1064.
- Phillips, N., Dennerstein, L., & Farish, S. (1996). Psychological morbidity in obstetric-gynaecology patients: testing the need for expanded psychiatry services in obstetric gynaecology facilities. *Australian and New Zealand Journal of Psychiatry*, 30, 74–81.
- Pigott, T. A. (1998). Obsessive-compulsive disorder: Symptom overview and epidemiology. *Bulletin of the Menninger Clinic*, 62(4, Suppl A), A4–A32.
- Pigott, T. A. (1999). Gender differences in the epidemiology and treatment of anxiety disorders. *Journal of Clinical Psychiatry*, 60(Suppl 18), 4–15.
- Pomerantz, E. M., & Ruble, D. N. (1998). The role of maternal gender socialization in the development of sex differences in child self-evaluative mechanisms. *Child Development*, 68, 558–478.
- Poulton, R., Milne, B. J., Craske, M. G., & Menzies, R. G. (2001). A longitudinal study of the etiology of separation anxiety. *Behaviour Research and Therapy*, 39(12), 1395–1410.
- Rasmussen, S. A., & Eisen, J. L. (1990). Epidemiology of obsessive compulsive disorder. *Journal of Clinical Psychiatry*, 51(2, Suppl), 10–13.
- Rauste-von Wright, M., von Wright, J., & Frankenhauser, M. (1981). Relationships between sex-related psychological characteristics during adolescence and catecholamine excretion during achievement stress. *Psychophysiology*, 18(4), 362–370.
- Reiss, S., & McNally, R. J. (1985). The expectancy model of fear. In S. Reiss & R. R. Bootzin (Eds.), *Theoretical issues in behavior therapy* (pp. 107–121). New York: Academic Press.
- Robichaud, M., Dugas, M. J., & Conway, M. (2003). Gender differences in worry and associated cognitive-behavioral variables. *Journal of Anxiety Disorders*, 17, 501–516.
- Rohrmann, S., Netter, P., Hennig, J., & Hodapp, W. (2003). Repression-sensitization, gender, and discrepancies in psychobiological reactions to examination stress. *Anxiety, Stress & Coping: An International Journal*, 16(3), 321–329.
- Ross, C. E., & Mirowsky, J. (2002). Age and the gender gap in the sense of personal control. *Social Psychology Quarterly*, 65(2), 125–145.
- Ross, L. E., & McLean, L. M. (2006). Anxiety disorders during pregnancy and the postpartum period: A systematic review. *Journal of Clinical Psychiatry*, 67(8), 1285–1298.
- Scheibe, G., & Albus, M. (1992). Age at onset, precipitating events, sex distribution, and co-occurrence of anxiety disorders. *Psychopathology*, 25(1), 11–18.
- Schmidt, N. B., & Kosselka, M. (2000). Gender differences in patients with panic disorder: Evaluating cognitive mediation of phobic avoidance. *Cognitive Therapy and Research*, 24(5), 533–550.
- Sheikh, J. I., Leskin, G. A., & Klein, D. F. (2002). Gender differences in panic disorder: Findings from the National Comorbidity survey. *American Journal of Psychiatry*, 159(1), 55–58.
- Silverman, W. K., La Greca, A. M., & Wasserstein, S. (1995). What do children worry about? Worries and their relationship to anxiety. *Child Development*, 66, 671–686.
- Skre, I., Onstad, S., Torgersen, S., Lygren, S., & Kringlen, E. (1993). A twin study of DSM-III-R anxiety disorders. *Acta Psychiatrica Scandinavica*, 88(2), 85–92.
- Spielberger, C. D. (1966). Theory and research on anxiety. In C. D. Spielberger (Ed.), *Anxiety and behavior* (pp. 3–19). New York: Academic Press.
- Spielberger, C. D. (1972). Conceptual and methodological issues in anxiety research. In C. D. Spielberger (Ed.), *Anxiety: Current trends in theory and research* (pp. 481–492). New York: Academic Press.
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. D. (1970). *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologists Press.
- Stein, M., Jang, K., & Livesley, W. (2002). Heritability of social anxiety-related concerns and personality characteristics: A twin study. *Journal of Nervous and Mental Disease*, 190(4), 219–224.
- Steiner, H., Ryst, E., Berkowitz, J., Gschwendt, M. A., & Koopman, C. (2002). Boys' and girls' responses to stress: Affect and heart rate during a speech task. *Journal of Adolescent Health*, 30(Suppl), 14–21.
- Stevenson-Hinde, J., & Shouldice, A. (1993). Wariness to strangers: A behavior systems perspective revisited. In K. H. Rubin & J. B. Asendorpf (Eds.), *Social withdrawal, inhibition, and shyness in childhood* (pp. 101–116). New York: Lawrence Erlbaum.
- Stewart, S. H., Taylor, S., & Baker, J. M. (1997). Gender differences in dimensions of anxiety sensitivity. *Journal of Anxiety Disorders*, 11, 179–200.
- Ströhle, A. (2000). Increased response to a putative panicogenic nocebo administration in female patients with panic disorder. *Journal of Psychiatric Research*, 34(6), 439–442.
- Stoney, C. M., Davis, M. C., & Matthews, K. A. (1987). Sex differences in physiological responses to stress and in coronary heart disease: A causal link? *Psychophysiology*, 24(2), 127–131.
- Suarez, L., & Bell-Dolan, D. (2001). The relationship of child worry to cognitive biases: Threat interpretation and likelihood of event occurrence. *Behavior Therapy*, 32(3), 425–442.
- Taylor, S. (1998). The hierarchic structure of fears. *Behaviour Research and Therapy*, 36(2), 205–214.
- Taylor, S. E., Klein, L. C., Lewis, B. P., Gruenewald, T. L., Gurung, R. A. R., & Updegraff, J. A. (2000). Biobehavioral responses to stress in females: Tend-and-befriend, not fight-or-flight. *Psychological Review*, 107(3), 411–429.
- Telch, M. J., Lucas, J. A., & Nelson, P. (1989). Nonclinical panic in college students: An investigation of prevalence and symptomatology. *Journal of Abnormal Psychology*, 98(3), 300–306.
- Thoits, P. A. (1987). Gender and marital status differences in control and distress: Common stress versus unique stress explanations. *Journal of Health and Social Behavior*, 28, 7–22.
- Thorpe, S. J., & Salkovskis, P. M. (1995). Phobic beliefs: Do cognitive factors play a role in specific phobias? *Behaviour Research and Therapy*, 33, 805–816.
- Todman, J., & Day, K. (2006). Computer anxiety: The role of psychological gender. *Computers in Human Behavior*, 22(5), 856–869.
- Tolin, D. F., & Foa, E. B. (2006). Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychological Bulletin*, 132, 959–992.
- True, W. R., Rice, J., Eisen, S. A., Goldberg, J., Lyons, M. J., & Nowack, J. A. (1993). A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Archives of General Psychiatry*, 50(4), 257–264.
- Tucker, M., & Bond, N. W. (1997). The roles of gender, sex role, and disgust in fear of animals. *Personality and Individual Differences*, 22(1), 135–138.
- Turgeon, L., Marchand, A., & Dupuis, G. (1998). Clinical features in panic disorder with agoraphobia: A comparison of men and women. *Journal of Anxiety Disorders*, 12(6), 539–553.
- Watt, M. C., Stewart, S. H., & Cox, B. J. (1998). A retrospective study of the learning origins of anxiety sensitivity. *Behaviour Research and Therapy*, 36, 05–525.
- Weinberg, M. K., Tronick, E. Z., Cohn, J. F., & Olson, K. L. (1999). Gender differences in emotional expressivity and self-regulation during early infancy. *Developmental Psychology*, 35(1), 175–188.
- Weissman, M. M., & Merikangas, K. R. (1986). The epidemiology of anxiety and panic disorders: An update. *Journal of Clinical Psychiatry*, 47(Suppl), 11–17.
- Wheaton, M., Timpano, K. R., LaSalle-Ricci, V. H., & Murphy, D. (2008). Characterizing the hoarding phenotype in individuals with OCD: Associations with comorbidity, severity and gender. *Journal of Anxiety Disorders*, 22, 243–252.
- Williams, K. E., & Koran, L. M. (1997). Obsessive-compulsive disorder in pregnancy, the puerperium, and the premenstruum. *Journal of Clinical Psychiatry*, 58, 330–334.
- Wood, W., & Eagly, A. H. (2002). A cross-cultural analysis of the behavior of women and men: Implications for the origins of sex difference. *Psychological Bulletin*, 128, 699–727.
- Wolpe, J., & Lang, P. J. (1977). *Manual for the Fear Survey Schedule*. San Diego, CA: Educational and Industrial Testing Service.
- Yonkers, K. A., Zlotnick, C., Allsworth, J., Warshaw, M., Shea, T., & Keller, M. B. (1998). Is the course of panic disorder the same in women and men? *American Journal of Psychiatry*, 155(5), 596–602.
- Zinbarg, R. E., & Barlow, D. H. (1996). Structure of anxiety and the anxiety disorders: A hierarchical model. *Journal of Abnormal Psychology*, 105(2), 181–193.
- Zinbarg, R. E., Barlow, D. H., & Brown, T. A. (1997). Hierarchical structure and general factor structure saturation of the Anxiety Sensitivity Index: Evidence and implications. *Psychological Assessment*, 9, 277–284.
- Zvolensky, M. J., McNeil, D. W., Porter, C. A., & Stewart, S. H. (2001). Assessment of anxiety sensitivity in young American Indians and Alaska Natives. *Behaviour Research and Therapy*, 39, 477–493.